

## $\beta$ -Alanine supplementation augments muscle carnosine content and attenuates fatigue during repeated isokinetic contraction bouts in trained sprinters

Wim Derave,<sup>1</sup> Mahir S. Özdemir,<sup>2</sup> Roger C. Harris,<sup>3</sup> Andries Pottier,<sup>1</sup> Harmen Reyngoudt,<sup>4</sup> Katrien Koppo,<sup>1</sup> John A. Wise,<sup>5</sup> and Eric Achten<sup>4</sup>

<sup>1</sup>Department of Movement and Sport Sciences, and <sup>2</sup>Department of Electronics and Information Systems, Medical Image and Signal Processing, Ghent University, Ghent, Belgium; <sup>3</sup>School of Sports, Exercise and Health Sciences, University of Chichester, Chichester, United Kingdom; <sup>4</sup>Department of Radiology, Ghent Institute for Functional Magnetic Resonance Imaging, Ghent University, Ghent, Belgium; and <sup>5</sup>Natural Alternatives International, San Marcos, California

Submitted 13 April 2007; accepted in final form 2 August 2007

**Derave W, Özdemir MS, Harris RC, Pottier A, Reyngoudt H, Koppo K, Wise JA, Achten E.**  $\beta$ -Alanine supplementation augments muscle carnosine content and attenuates fatigue during repeated isokinetic contraction bouts in trained sprinters. *J Appl Physiol* 103: 1736–1743, 2007. First published August 9, 2007; doi:10.1152/jappphysiol.00397.2007.—Carnosine ( $\beta$ -alanyl-L-histidine) is present in high concentrations in human skeletal muscle. The ingestion of  $\beta$ -alanine, the rate-limiting precursor of carnosine, has been shown to elevate the muscle carnosine content. We aimed to investigate, using proton magnetic resonance spectroscopy (proton MRS), whether oral supplementation with  $\beta$ -alanine during 4 wk would elevate the calf muscle carnosine content and affect exercise performance in 400-m sprint-trained competitive athletes. Fifteen male athletes participated in a placebo-controlled, double-blind study and were supplemented orally for 4 wk with either 4.8 g/day  $\beta$ -alanine or placebo. Muscle carnosine concentration was quantified in soleus and gastrocnemius by proton MRS. Performance was evaluated by isokinetic testing during five bouts of 30 maximal voluntary knee extensions, by endurance during isometric contraction at 45% maximal voluntary contraction, and by the indoor 400-m running time.  $\beta$ -Alanine supplementation significantly increased the carnosine content in both the soleus (+47%) and gastrocnemius (+37%). In placebo, carnosine remained stable in soleus, while a small and significant increase of +16% occurred in gastrocnemius. Dynamic knee extension torque during the fourth and fifth bout was significantly improved with  $\beta$ -alanine but not with placebo. Isometric endurance and 400-m race time were not affected by treatment. In conclusion, 1) proton MRS can be used to noninvasively quantify human muscle carnosine content; 2) muscle carnosine is increased by oral  $\beta$ -alanine supplementation in sprint-trained athletes; 3) carnosine loading slightly but significantly attenuated fatigue in repeated bouts of exhaustive dynamic contractions; and 4) the increase in muscle carnosine did not improve isometric endurance or 400-m race time.

buffer capacity; ergogenic supplements; nuclear magnetic resonance; exercise performance; track and field

CARNOSINE ( $\beta$ -alanyl-L-histidine) is a cytoplasmic dipeptide synthesized from the precursors L-histidine and  $\beta$ -alanine by carnosine synthetase. Carnosine is present in relatively high concentrations in skeletal muscle (5–10 mM) and other excitable tissues, such as nervous tissue. Given that the  $pK_a$  value of the imidazole ring is 6.83 (4, 42), carnosine acts as a physicochemical buffer in myocytes. This function is consistent with the fact that carnosine is present in higher concentrations in glycolytic than in oxidative muscle fibers (7, 8), with the

highest levels found in animals that perform frequent sprint exercises (e.g., greyhounds, thoroughbred horses), explosive flight behaviors (e.g., pheasants), and prolonged hypoxic dives (e.g., whales) (16, 40). In addition, it has been shown that carnosine causes an increase in the  $Ca^{2+}$  sensitivity of vertebrate skeletal muscle. Given that the evidence for the latter is solely based on in vitro studies with skinned muscle fibers of frogs and rats (11, 26), its physiological relevance in vivo remains to be established.

Several studies have demonstrated, by a preexercise metabolic alkalosis intervention, the importance of intramyocellular pH regulation in performance during supramaximal exercise (29), probably via an indirect effect of proton accumulation on contractile function. While carnosine will function as a buffer over the physiological pH range and has been estimated to account for ~10% of the total buffering capacity in human vastus lateralis (19), the importance of buffering and pH control in muscle to exercise performance is still debated. By using a newly developed nutritional strategy to raise carnosine concentrations in human muscle, resulting in altered buffering capacity (18, 19), there is now a means to undertake studies equating muscle buffering capacity to performance. In Harris et al. (18), a daily oral intake of ~6 g of  $\beta$ -alanine for 4 wk was shown to increase the carnosine content by 60% in the vastus lateralis of untrained humans, while 10-wk supplementation increased this by ~80% (19). The pronounced effect of  $\beta$ -alanine supplementation is consistent with evidence that the intramyocellular synthesis of carnosine is rate limited by the availability of  $\beta$ -alanine (3, 9, 19) and not by histidine, which is present in muscle in high concentrations relative to its  $K_m$  for carnosine synthase (20). Thus the use of  $\beta$ -alanine supplementation offers researchers a technique to investigate the role of carnosine in muscle contraction, as well as its ergogenic potential.

Several recent studies have suggested that muscle carnosine content is a determining factor in high-intensity dynamic and isometric exercise performance (19, 39, 41). Suzuki and colleagues (41) observed a positive correlation between the vastus lateralis carnosine content and the power output generated at the end of a 30-s all-out cycling sprint (Wingate-test) in untrained men. Additionally, Hill et al. (19) supplemented untrained men with  $\beta$ -alanine for 10 wk, which resulted not only in a substantial increase in the carnosine content of the

Address for reprint requests and other correspondence: W. Derave, Dept. of Movement and Sport Sciences, Ghent Univ., Watersportlaan 2, B-9000 Ghent, Belgium (e-mail: Wim.derave@ugent.be).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

vastus lateralis, but also in an increased time to exhaustion in a cycling test performed at 110% of power output at maximal heart rate, where the expected endurance time is  $\sim 2.5$  min. In the present study, the ergogenic potential of supplementary  $\beta$ -alanine is evaluated in sprint-trained subjects, in which there is added interest because sprint-trained athletes are reported to have a higher muscle carnosine content than untrained subjects or endurance-trained athletes (31). A carnosine content that is initially high could mitigate against the effectiveness of  $\beta$ -alanine supplementation to further elevate muscle carnosine. By analogy, the latter "ceiling" phenomenon is known to limit the effectiveness of creatine supplementation in subjects with high initial muscle creatine content (13, 17).

For the quantification of the human muscle carnosine concentrations, there are currently only two techniques used, i.e., the determination of imidazole dipeptides by HPLC or by ion-exchange chromatography of extracts of muscle. Both of these require muscle biopsy samples (8, 41). However, proton magnetic resonance spectroscopy (MRS) could provide a valuable noninvasive alternative (30, 36). The unique properties of the imidazole protons of the histidine subunit of carnosine make them visible in the far downfield region [7 and 8 parts/million (ppm); left from the water peak] of the proton spectrum of human muscle (30). A second aim of this study was, therefore, to explore the possibility to quantify the carnosine content in the soleus and gastrocnemius muscles of the human calf through proton MRS.

The present study was conducted on well-trained, 400-m sprint runners. We hypothesized that muscle carnosine content would be substantially elevated by 4 wk of oral  $\beta$ -alanine supplementation, despite the possibility of a high initial carnosine content in this population. Because carnosine is known to act as a proton buffer in the physiological pH range, and because intramyocellular acidosis has been proposed as a cause of fatigue during high-intensity contractions, we further aimed to investigate the effect of the potentially elevated muscle carnosine content on several aspects of muscle performance and fatigue. This was investigated through 1) an isokinetic test of repeated bouts of maximal exhaustive knee extensions, 2) an isometric test of the static endurance of the knee extensors, and 3) an indoor 400-m race. Finally, we aimed to evaluate the use of proton MRS as a noninvasive tool to monitor muscle carnosine concentration in humans.

## MATERIALS AND METHODS

**Subjects.** Fifteen male track-and-field athletes, each with a personal record on 400 m below 53 s, were recruited from Flemish athletics clubs. Their official personal record on 400 m was, on average ( $\pm$ SD),  $50.45 \pm 1.60$  s and ranged between 47.49 and 52.64 s. All athletes were actively competing at regional or national (Belgium) level and were preparing for the indoor competition season during the course of the study (months of November and December). In the randomization, the number of athletes from the same club and trainer was matched for each group to minimize the differences in training amount and intensity. The athletes of the placebo ( $n = 7$ ) and  $\beta$ -alanine ( $n = 8$ ) groups performed, on average, 5.4 and 5.6 training sessions per week, respectively, during the study period. The relative percentage of total training time spent on power, speed, resistance, and endurance was 25, 23, 24, and 28 in placebo and 27, 25, 23, and 25 in  $\beta$ -alanine, respectively. None of the subjects was taking any other oral ergogenic supplements during the study or had taken supplements in the 3 mo before the study. The subjects gave their informed consent, and the

study was approved by the local Ethics Committee (Ghent University Hospital, Belgium). The subjects' weight, height, and age were  $70.7 \pm 5.7$  kg,  $183 \pm 4$  cm, and  $18.4 \pm 1.5$  yr for placebo and  $74.1 \pm 7.2$  kg,  $184 \pm 8$  cm, and  $23.8 \pm 4.2$  yr for  $\beta$ -alanine, respectively. All subjects completed all experiments, and there were no complaints of side-effects of the supplements.

**Study design.** A placebo-controlled, double-blind study was performed. Subjects were supplemented orally for 4–5 wk with either placebo (maltodextrin) or  $\beta$ -alanine (Carnosyn, National Alternatives International, San Marcos, CA). Supplements were provided in capsules of 400 mg and were administered each day as six divided doses, with at least 2 h in between ingestions. Daily doses consisted of 2.4 g/day during the first 4 days, 3.6 g/day during the subsequent 4 days, and from then on 4.8 g/day until the end of the study. All measurements were performed in the week before the start of the supplementation (Pre) and during the last week of supplementation (Post). As for the muscle torque measurements, a familiarization session was performed 1 wk before the Pre measurements.

**Proton MRS.** All of the MRS measurements were performed on a 3-Tesla whole body MRI scanner (Siemens Trio, Erlangen) equipped with a knee coil. Single voxel point-resolved spectroscopy with the following parameters was used: repetition time (TR) = 2,000 ms, echo time (TE) = 30 ms, voxel size = 40 mm  $\times$  30 mm  $\times$  12 mm, number of excitation = 256, 1,024 data points, spectral bandwidth of 1,200 Hz, and a total acquisition time of 8.40 min. A 500-ml spherical container filled with an aqueous solution of 50 mM carnosine (Sigma) was used as an external reference for the absolute quantification. The full-width-half maximum of the water signal was minimized to be in the range of 20–30 Hz during shimming.

The following equation was used to determine the concentration of C2-H carnosine in vivo:

$$C_m = \frac{C_r S_m F_{mT_1} F_{mT_2}}{S_r F_{rT_1} F_{rT_2}} \quad (1)$$

where  $C_m$  is the carnosine concentration in vivo;  $C_r$  is the concentration of the external reference phantom prepared (50 mM);  $F_{mT_1}$  and  $F_{mT_2}$  are correction factors for the  $T_1$  and  $T_2$  relaxation of the metabolites of interest, respectively; and  $S_m$  and  $S_r$  are the estimated signal peak areas obtained by the curve fitting performed in the frequency domain. Data processing included 5-Hz apodization (line broadening), phase correction, baseline correction, and Fourier transformation. The correction factors  $F_{T_1}$  and  $F_{T_2}$  were calculated as follows:

$$F_{T_1} = \frac{1}{1 - \exp(-TR/T_1)} \quad (2)$$

$$F_{T_2} = \frac{1}{\exp(-TE/T_2)} \quad (3)$$

$T_1$  and  $T_2$  relaxation parameters of in vitro carnosine were measured and were found to be  $560 \pm 15$  and  $66.22 \pm 9.3$  ms, respectively. Thus, for the pulse sequence used (TR = 2,000 ms), the correction term in Eq. 2 amounts to 0.9718 and can therefore be neglected. Note that this correction factor for in vivo carnosine would be larger than that of in vitro due to shorter  $T_1$  of the carnosine in vivo.  $T_2$  in vivo used in correction was 39 ms, as was previously measured (30). Signal intensities were also corrected for the differences in coil loading and temperature difference between the phantom and the human body. Load correction was carried out using the following formula:

$$S_c = S_u 10^{A/20} \quad (4)$$

where  $S_c$  is the corrected and  $S_u$  is the uncorrected signal intensities, and  $A$  is the difference between radio frequency power required for 180° pulse in vivo and that for phantom, measured in decibels (38). The correction factor used for the temperature difference was 6% (44).

**Isokinetic and isometric muscle fatigue protocol.** An isokinetic and isometric test was performed on a Biodex isokinetic dynamometer (System 3, Biodex Medical Systems) to evaluate the contractile performance of the knee extensors. The dynamometry was preceded by a standardized warm-up consisting of an ascent and descent of a stairs (11-m altitude) and two 50-m runs at moderate pace. The isokinetic test was performed on the right leg. This consisted of  $5 \times 30$  maximal voluntary isokinetic knee extensions at a constant angular velocity of  $180^\circ/\text{s}$ . Each contraction was initiated from a position of  $90^\circ$  knee flexion and was continued to the point of full knee extension. After each extension, the lower leg was passively returned to the start position at  $90^\circ/\text{s}$  ( $\sim 1.5$  s for full cycle). Each bout of 30 contractions was separated by a 1-min recovery period. Subjects were encouraged during the first three contractions to make sure that they were contracting maximally from the start of each bout. Subjects received visual feedback of their produced peak torque. Peak torque during each contraction (1–30) was measured and used to calculate the average peak torque during each bout of exercise. Subsequently, the isometric test was performed on the left leg. The knee angle was fixed at  $45^\circ$ , and the maximal static voluntary contraction (MVC) torque was determined. The highest torque of three 3-s attempts, separated by 30 s of rest, was considered as MVC. Subjects were then asked to contract isometrically with their knee extensors at a target torque of 45% of MVC for as long as possible to determine isometric endurance.

**400-m race.** The time to complete 400-m running was evaluated in an indoor 300-m flat athletics track (Flanders Sports Arena, Ghent, Belgium). Time was recorded by an infrared light-based electronic device at the start and finish (400 m). Subjects were asked to warm up, according to their personal routine during  $\sim 45$  min. Following warm-up and 2–3 min before the start, a capillary blood sample was obtained from a finger tip and analyzed for lactate with Lactate PRO test strips (Arkray, Kyoto, Japan). Also at 90 and 180 s following finish, blood lactate was measured. The maximal lactate accumulation was considered as the higher of the two postprint lactate values.

**Statistics.** A  $2 \times 2$  general linear model repeated-measures ANOVA was used to evaluate the muscle carnosine content, 400-m run time, and lactate accumulation and isometric endurance, with “group” (placebo vs.  $\beta$ -alanine) as between-subjects factor and time (Pre vs. Post) as a within-subjects factor (SPSS statistical software, SPSS, Chicago, IL). The isokinetic contraction bouts were evaluated with a  $2 \times 2 \times 5$  repeated-measures ANOVA, with “bout” as an additional within-subjects factor. In case of significant interaction, ANOVA was repeated at a lower level on each half of the data. Values are presented as means  $\pm$  SD, and significance was assumed at  $P < 0.05$ .

## RESULTS

**Muscle carnosine content.** Typical proton MRS spectra and voxel localization of the soleus and gastrocnemius muscle are shown in Fig. 1. In carnosine phantoms and in human soleus muscle, the C4-H and C2-H peaks have very similar size. However, in gastrocnemius, the C4-H peak was markedly and consistently smaller than the C2-H peak in all subjects (Fig. 1D), which appears to agree with the findings of others, publishing a spectrum from a human gastrocnemius muscle (37, 46). Our additional experiments revealed that the test-retest reproducibility was good for the C2-H but poor for C4-H peak (M. S. Özdemir, H. Reyngoudt, Y. De Deene, H. S. Sazak, E. Fieremans, S. Delputte, Y. D’Asseler, W. Derave, I. Lemahieu, and E. Achten, unpublished observations). This high variation of the C4 resonance and the low C4-to-C2 ratio in gastrocnemius both suggest marked orientation-dependent effects on this resonance (24) and led to the current C2-




Fig. 1. Proton magnetic resonance spectroscopy (MRS) and MRI of the right calf muscles in a representative sprint-trained subject. Full proton spectra of the soleus (A) and gastrocnemius (B) are shown. Detailed spectra of the downfield carnosine region in soleus (C) and gastrocnemius (D) are shown. Resonances for the C2-H and C4-H imidazole protons of carnosine are at  $\sim 8$  and  $\sim 7$  parts/million (ppm), respectively. Coronal (E and F) and transverse (G and H) images were obtained by MRI in soleus (E and G) and gastrocnemius (F and H) to demonstrate the localization of the voxel.

based method for carnosine quantification. In the soleus muscle (Fig. 2A), the absolute carnosine concentration as measured by MRS before supplementation was  $7.25 \pm 1.47$  and  $7.76 \pm 1.36$  mmol/l in the placebo and  $\beta$ -alanine groups, respectively. As a result of supplementation, the carnosine content of the soleus increased by 47% ( $P < 0.0001$ ) in the  $\beta$ -alanine group to  $11.39 \pm 1.38$  mmol/l, whereas it remained stable at  $7.85 \pm 1.04$  mmol/l (+8%,  $P = 0.41$ ) in the placebo group. In the gastrocnemius, the baseline carnosine concentra-






Fig. 2. Muscle carnosine concentration (mmol/l) in soleus (A) and gastrocnemius (B) is shown before and after 4-wk supplementation of placebo or  $\beta$ -alanine. C: the individual change in muscle carnosine content [Post (last week of supplementation) vs. Pre (week before supplementation)] in the 7 placebo subjects and 8  $\beta$ -alanine subjects. The horizontal line depicts the average procentual change for each group. D: the absolute change in muscle carnosine concentration is shown in relation to the initial carnosine content of soleus ( $\circ$ ) and gastrocnemius ( $\bullet$ ) in each of the  $\beta$ -alanine-supplemented subjects. \*Different from Pre ( $P < 0.05$ ).<sup>o</sup>Different from placebo ( $P < 0.05$ ).

tion was  $\sim 25\%$  higher than in soleus ( $P < 0.05$ ), being  $8.56 \pm 1.88$  and  $10.16 \pm 1.91$  mmol/l in the placebo and  $\beta$ -alanine groups, respectively. The gastrocnemius carnosine content (Fig. 2B) increased significantly in both groups over the 4-wk supplementation period, but the increase was significantly more pronounced in the  $\beta$ -alanine group ( $+37\%$ ;  $P < 0.0001$ ) than in the placebo group ( $+16\%$ ;  $P = 0.005$ ). In subjects receiving  $\beta$ -alanine, the carnosine content of the soleus increased by at least 28% and maximally 89% and in the gastrocnemius between 22 and 53% (see Fig. 2C). As shown in Fig. 2D, the increase in carnosine in the  $\beta$ -alanine-supplemented subjects was not dependent on the initial muscle carnosine content. The lack of negative correlation ( $r = +0.18$ ) suggests that there was no ceiling effect and that a maximal carnosine content may not have been attained. The highest carnosine concentration measured was 18.5 mM observed in the gastrocnemius muscle of a subject given  $\beta$ -alanine supplementation, which was a 40% increase over baseline.

**Repeated bouts of maximal isokinetic knee extensions.** The average knee extension torque produced during five bouts of 30 dynamic (isokinetic) contractions is shown in Table 1. We performed a  $2 \times 2 \times 5$  ANOVA (with "group", "Pre/Post", and "bout" as factors) on these data, and we found a significant three-way interaction ( $P = 0.05$ ). Therefore, the data were further analyzed with two-way ANOVAs in both groups separately. In the placebo group, a significant Pre/Post  $\times$  bout

interaction was found, and post hoc analysis revealed a significant Pre/Post effect in the first two bouts, but not in the subsequent three bouts. In the  $\beta$ -alanine group, however, a significant Pre/Post main effect was observed, indicating that average torque was higher in all five bouts Post vs. Pre supplementation. This effect is further illustrated in more detail in Fig. 3. The peak knee extension torque of each contraction during *bouts 4* and *5* is consistently higher Post  $\beta$ -alanine supplementation compared with Pre ( $+6.1\%$  in *bout 4* and  $+3.8\%$  in *bout 5*), whereas in the placebo group the Post and Pre values overlap ( $+1.0\%$  in *bout 4* and  $-0.8\%$  in *bout 5*).

**Isometric endurance.** The maximal duration of a sustained isometric (static) knee extensor contraction at 45% MVC was  $173 \pm 55$  and  $201 \pm 48$  s at baseline and increased to  $187 \pm 64$  and  $220 \pm 61$  s following supplementation of placebo or  $\beta$ -alanine, respectively. ANOVA revealed a significant Pre/Post main effect ( $P = 0.03$ ), but no significant group main effect or interaction ( $P = 0.66$ ).

**400-m race.** The mean time to complete a 400-m run on the indoor athletic track was  $52.17 \pm 1.83$  and  $51.11 \pm 1.66$  s at baseline for placebo and  $\beta$ -alanine subjects, respectively. Times decreased on average by  $\sim 0.7$  s to  $51.44 \pm 1.57$  and  $50.36 \pm 1.43$  s, respectively, following supplementation in both groups. The improved run time was significant but similar in both groups, as illustrated by the main Pre/Post effect ( $P = 0.02$ ) and the lack of Pre/Post  $\times$  group interaction ( $P = 0.98$ ).

Table 1. Average knee extension torque during 5 bouts of 30 dynamic contractions

	Bout 1	Bout 2	Bout 3	Bout 4	Bout 5	Pre vs. Post	Bout	Interaction
<i>Placebo</i>								
Pre	167 (23)	150 (15)	143 (19)	142 (20)	144 (18)	0.169	<0.001	0.010
Post	178 (21)*	158 (16)*	148 (14)	143 (13)	143 (13)			
<i><math>\beta</math>-Alanine</i>								
Pre	176 (22)	148 (19)	137 (18)	132 (20)	133 (21)	0.011	<0.001	0.739
Post	181 (25)*	155 (23)*	143 (22)*	139 (23)*	137 (25)*			

Values are means (SD) in Nm. Pre, week before supplementation; Post, last week of supplementation. The  $P$  values for the main effects (Pre/Post and bout) and for the interaction of the  $2 \times 5$  ANOVA are shown and underlined in the case of significance. \*Significant difference vs. Pre.




Fig. 3. Peak knee extension torque during each contraction of *bout 4* (A and C) and *bout 5* (B and D) before and after 4 wk of placebo (A and B) or  $\beta$ -alanine (C and D) supplementation. Shown are the average of 7 (placebo) and 8 ( $\beta$ -alanine) subjects. E: delta (Post vs. Pre) average torque of all contractions in one bout (means  $\pm$  SE). \*Difference between Post and Pre ( $P < 0.05$ ).

in a  $2 \times 2$  ANOVA. The individual changes in run time are shown in Fig. 4, A (placebo) and B ( $\beta$ -alanine). Similarly, the blood lactate accumulation 180 s following the 400-m sprint was significantly increased by  $\sim 1.2$  mmol/l in both groups following supplementation (Table 2). The individual changes in maximal blood lactate concentrations following the 400-m are shown in Fig. 4, C (placebo) and D ( $\beta$ -alanine). The changes in gastrocnemius carnosine content did not significantly correlate with changes in 400-m running speed ( $r = 0.08$ ), in maximal lactate accumulation ( $r = -0.22$ ), or with isometric endurance ( $r = 0.12$ ).

## DISCUSSION

The present data on muscle carnosine concentration, obtained by proton MRS, agree closely with data from muscle biopsy studies of the vastus lateralis, showing that the muscle carnosine content can be substantially increased by  $\beta$ -alanine supplementation (18, 19). The subjects in this study, however, were sprint-trained competitive athletes rather than untrained subjects, and the muscles studied were calf muscles (soleus and gastrocnemius) rather than thigh muscles (vastus lateralis). The magnitude of the increase in muscle carnosine content induced by 4.8 g/day  $\beta$ -alanine supplementation in the present study, namely +47% in soleus and +37% in gastrocnemius, is comparable yet slightly lower than in a previous report on untrained subjects (+64% in vastus lateralis) following 4 wk of 4 g/day, increasing to 6.4 g/day, of  $\beta$ -alanine supplementation (19).

As illustrated in Fig. 2D, even the subjects with a very high initial carnosine content ( $>12$  mmol/l) were still able to elevate their concentrations by an additional 4–5 mmol/l. This suggests a lack of any “ceiling effect” and that the human muscle carnosine content is far from maximal. Furthermore, this is supported by the finding that, following 4 wk of  $\beta$ -alanine supplementation, an additional 6 wk of supplementation caused an additional 20% increase in muscle carnosine (19). In this respect, carnosine is clearly different from creatine, because maximal muscle creatine content is already obtained after 1 wk of oral creatine supplementation, and subjects with high initial levels are often nonresponders (6, 17).

A further aim of this study was to evaluate the potential ergogenic effect of muscle carnosine loading in trained sprinters. The results show that, while elevated muscle carnosine content leads to an improvement in muscle torque during repeated bouts of intense dynamic contractions, it did not apparently affect 400-m running performance. It has long been suggested that the acidotic environment that is created by carboxylate anion formation during anaerobic work limits muscle contractility (5), and there is a vast amount of literature suggesting that preexercise alkalosis promotes performance in brief, high-intensity exercise (27, 34). Several papers with a similar study population (runners) and exercise type (running race between 300- and 800-m distance), as with the present study, have investigated the effect of induced metabolic alkalosis through oral preexercise ingestion of sodium bicarbonate or sodium citrate. The outcome of these studies, however, has




Fig. 4. Time (s) to complete a 400-m run before and after 4 wk of placebo (A) or  $\beta$ -alanine (B) supplementation is shown. Maximal capillary blood lactate concentration (mmol/l) following a 400-m run before and after 4 wk of placebo (C) or  $\beta$ -alanine (D) supplementation is shown. Individual values (shaded symbols) and group averages (open symbols) are shown.

been equivocal, with two studies reporting a positive effect (12, 45) and other studies reporting no effect (21, 23, 43). In the present study, the increase in muscle carnosine content in the  $\beta$ -alanine-supplemented group will inevitably give rise to an increase in the intramyocellular buffer capacity. The lack of difference (interaction term of 0.98) in 400-m race time between placebo and  $\beta$ -alanine would indicate either that the gain in intracellular buffer capacity in this case was insufficient to bring about an increase in performance over and above the limitations of the test, or that decreasing pH is not a limiting factor to 400-m performance in trained individuals. The apparent contradiction with earlier data by Goldfinch et al. (12), reporting a 1.5-s improvement in 400-m race time following bicarbonate ingestion, may in part relate to differences in training status, since the male subjects in the Goldfinch study

had average baseline 400-m times of 58.5 s (74% of current world record speed), compared with 51.6 s (84% of current world record speed) in the present study. Moreover, we must be cautious when interpreting the negative outcome of the performance part of the present study. As it takes several weeks of supplementation to induce muscle carnosine loading and because the washout period is unknown, it was not possible to adopt a paired crossover design. In an unpaired experimental design, the study population of seven to eight subjects per group possibly did not generate enough statistical power. Thus presently we cannot exclude that still higher elevations of muscle carnosine content (e.g., by using a longer supplementation period) or a study with a more powerful statistical design may still demonstrate an increase in sprint performance with  $\beta$ -alanine supplementation.

Table 2. Capillary blood lactate accumulation before the start (post-warm-up) and 90 and 180 s after the finish of a 400-m run

Lactate	Pre	Post	Pre/Post	Group	Interaction
Post-warm-up			0.208	0.554	0.457
Placebo	2.5 (0.9)	3.4 (1.4)			
$\beta$ -alanine	3.1 (1.1)	3.3 (0.9)			
90 s Post-finish			0.162	0.248	0.723
Placebo	14.7 (1.8)	15.4 (1.4)			
$\beta$ -alanine	15.5 (0.9)	16.0 (1.2)			
180 s Post-finish			<u>0.003</u>	0.441	0.989
Placebo	14.7 (1.3)	15.9 (0.7)*			
$\beta$ -alanine	15.1 (1.4)	16.3 (0.8)*			

Values are means (SD) in mmol/l. The *P* values for the main effects (Pre/Post and group) and for the interaction of the  $2 \times 2$  ANOVA are shown and underlined in the case of significance. \*Significant difference vs. Pre.

Dynamic knee extension torque was significantly increased in each of the five bouts of 30 maximal contractions by  $\beta$ -alanine supplementation, whereas, in the placebo group, only the first two bouts were improved. These data for the first time show that, in trained athletes performing repeated exhaustive contractions, muscle fatigue can be slightly but significantly attenuated in the later (*bouts 4 and 5*) stages of exercise by carnosine loading. These data markedly resemble those of the first study to address the effect of muscle creatine loading on muscle fatigue (14). In a nearly identical protocol, muscle torque was found to be increased following creatine supplementation in most of the  $5 \times 30$  maximal contractions (14).

Although still debated (25), most researchers agree that the rise in extracellular  $K^+$  and the intramyocellular accumulation of  $P_i$ , inorganic ADP, and protons ( $H^+$ ) are important initiating factors of fatigue, with reduced sarcoplasmic reticulum  $Ca^{2+}$  release and reduced myofibrillar  $Ca^{2+}$  sensitivity as important consequences. The effects are ultimately a reduction in power output (2). Carnosine does not affect sarcoplasmic reticulum  $Ca^{2+}$  release (11), but has been shown *in vitro* to increase myofibrillar  $Ca^{2+}$  sensitivity, particularly in fast-twitch fibers (11, 26). Furthermore, carnosine may attenuate fatigue by its physicochemical buffering capacity, thereby alleviating the detrimental effects that intracellular  $H^+$  accumulation may have, as recently shown (29, 33).

However, despite the effect of  $\beta$ -alanine supplementation on force production during repeated dynamic contractions, no effect on isometric endurance was observed. The lack of any effect is in contrast to the preliminary report of the findings of Ponte et al. (32), where again endurance at 45% MVC was studied. However, in the study of Ponte et al., contractions of the knee extensors were performed with a knee angle of  $90^\circ$  (compared to  $45^\circ$  in the present study), resulting in endurance times close to the 80-s predicted by the Rohmert curve for 45% MVC (1, 35), where the circulation is essentially occluded. With this model, peak accumulation of carboxylate anion (principally lactate and pyruvate), measuring  $H^+$  formation, occurs at 45% MVC and at a rate of  $\sim 1.2 \text{ mmol}\cdot\text{kg dry muscle}^{-1}\cdot\text{s}^{-1}$  ( $0.4 \text{ mmol}\cdot\text{l}^{-1}\cdot\text{s}^{-1}$ ) (1). The longer mean endurance times observed in the present study suggest a lesser degree of circulatory impairment with the model used, a lower rate of carboxylate anion accumulation, and a smaller fall in pH. This would limit the possibility of any increase in carnosine exerting a measurable effect on endurance and could explain the difference in findings. Alternatively, the added effect of muscle carnosine loading in this case is relatively small, because the sprint-trained subjects already have a high initial carnosine content and endogenous muscle buffer capacity.

The present study is possibly the first to apply the technique of proton MRS to noninvasively quantify and monitor the absolute carnosine concentrations in human skeletal muscle with  $\beta$ -alanine supplementation. Others have shown that the downfield C2-H and C4-H imidazole resonances of the proton spectrum of human muscle belong to the histidine subunit of carnosine ( $\beta$ -alanyl-L-histidine) (30, 36, 46). Pan et al. (30) indirectly estimated the absolute carnosine concentration in the forearm muscle of two volunteers from comparison with the estimated creatine concentration and reported values of 5.13 and 8.06 mM. In the present study, the absolute quantification of the muscle carnosine content was obtained by

relating the C2-H imidazole peak at 8 ppm (Fig. 1) of the proton spectrum of the muscle with that of a phantom (50 mM model solution of carnosine). The resulting absolute concentrations are  $\sim 7 \text{ mM}$  for soleus and  $\sim 9 \text{ mM}$  for gastrocnemius at baseline in this population.

The carnosine content was 25% higher in gastrocnemius compared with soleus. These data agree with previous studies, where higher carnosine levels were found in rat gastrocnemius vs. soleus (28) and in fast-twitch vs. slow-twitch fibers of the camel (10), horse (7), and human (15, 19, 22). When adopting a probable fiber-type distribution (%type I/II) of 88/12 in soleus and 47/53 in gastrocnemius (19) and a type I/II carnosine concentration ratio of 0.60, we calculated that carnosine content should be 25% higher in gastrocnemius compared with soleus, which is exactly the same as what we actually measured. Thus the difference in carnosine content between the two muscles is probably entirely explained by the difference in their fiber-type composition. This observation may be promising for the future development of MRS-based techniques to noninvasively determine muscle fiber-type composition and fiber compartmentalization in humans. We believe that it is correct to state that we provide physiological validation of the use of proton MRS in the quantification of human muscle carnosine concentrations, based on the acceptable absolute values of carnosine concentrations, the magnitude of  $\beta$ -alanine-induced increase in carnosine content, and the difference in carnosine content between muscle types found in this study.

It can be concluded from this study that 1) proton MRS can be used to noninvasively quantify human muscle carnosine content; 2) muscle carnosine content can be substantially elevated by oral  $\beta$ -alanine supplementation in sprint-trained athletes; 3) muscle carnosine loading slightly but significantly attenuates fatigue in repeated bouts of exhaustive dynamic contractions; and 4) 4 wk of supplementation with  $\beta$ -alanine at the prescribed dose used in this study do not result in a significant improvement in 400-m run performance. This latter result may be a consequence of a too little increase in intramyocellular pH buffering capacity for an improvement to be detected within the limits of the experimental procedures used, or alternatively may indicate that, in trained athletes, 400-m running performance is not limited by intracellular pH decrease.

#### ACKNOWLEDGMENTS

The contribution of Audrey Baguet, Sofie Debaere, Dieter De Bie, and Davy Van Tricht is greatly acknowledged.

#### REFERENCES

- Ahlborg B, Bergstrom J, Ekelund LG, Guarnieri G, Harris RC, Hultman E, Nordesjo LO. Muscle metabolism during isometric exercise performed at constant force. *J Appl Physiol* 33: 224–228, 1972.
- Allen DG, Lannergren J, Westerblad H. Muscle cell function during prolonged activity: cellular mechanisms of fatigue. *Exp Physiol* 80: 497–527, 1995.
- Bakardjiev A, Bauer K. Transport of beta-alanine and biosynthesis of carnosine by skeletal muscle cells in primary culture. *Eur J Biochem* 225: 617–623, 1994.
- Bate Smith EC. The buffering of muscle in rigor; protein, phosphate and carnosine. *J Physiol (Lond)* 92: 336–343, 1938.
- Dennig H, Talbott JH, Edwards HT, Dill DB. Effect of acidosis and alkalosis upon capacity for work. *J Clin Invest* 9: 601–613, 1931.
- Derave W, Eijnde BO, Hespel P. Creatine supplementation in health and disease: what is the evidence for long-term efficacy? *Mol Cell Biochem* 244: 49–55, 2003.
- Dunnett M, Harris RC. Carnosine and taurine contents of different fibre types in the middle gluteal muscle of the Thoroughbred horse. *Equine Vet J* 18, Suppl: 214–217, 1995.



8. **Dunnett M, Harris RC.** High-performance liquid chromatographic determination of imidazole dipeptides, histidine, 1-methylhistidine and 3-methylhistidine in equine and camel muscle and individual muscle fibres. *J Chromatogr B Biomed Appl* 688: 47–55, 1997.
9. **Dunnett M, Harris RC.** Influence of oral  $\beta$ -alanine and histidine supplementation on the carnosine content of the gluteus medius. *Equine Vet J* 30, Suppl: 499–504, 1999.
10. **Dunnett M, Harris RC, Soliman MZ, Suwar AA.** Carnosine, anserine and taurine contents in individual fibres from the middle gluteal muscle of the camel. *Res Vet Sci* 62: 213–216, 1997.
11. **Dutka TL, Lamb GD.** Effect of carnosine on excitation-contraction coupling in mechanically-skinned rat skeletal muscle. *J Muscle Res Cell Motil* 25: 203–213, 2004.
12. **Goldfinch J, Mc Naughton L, Davies P.** Induced metabolic alkalosis and its effects on 400-m racing time. *Eur J Appl Physiol* 57: 45–48, 1988.
13. **Greenhaff PL, Bodin K, Soderlund K, Hultman E.** Effect of oral creatine supplementation on skeletal muscle phosphocreatine resynthesis. *Am J Physiol Endocrinol Metab* 266: E725–E730, 1994.
14. **Greenhaff PL, Casey A, Short AH, Harris R, Söderlund K.** Influence of oral creatine supplementation on muscle torque during repeated bouts of maximal voluntary exercise in man. *Clin Sci (Lond)* 84: 565–571, 1993.
15. **Harris RC, Dunnett M, Greenhaff PL.** Carnosine and taurine contents in individual fibres of human vastus lateralis muscle. *J Sports Sci* 16: 639–643, 1998.
16. **Harris RC, Marlin DJ, Dunnett M, Snow DH, Hultman E.** Muscle buffering capacity and dipeptide content in the Thoroughbred horse, greyhound dog and man. *Comp Biochem Physiol A* 97: 249–251, 1990.
17. **Harris RC, Söderlund K, Hultman E.** Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clin Sci (Lond)* 83: 367–374, 1992.
18. **Harris RC, Tallon MJ, Dunnett M, Boobis L, Coakley J, Kim HJ, Fallowfield JL, Hill CA, Sale C, Wise JA.** The absorption of orally supplied beta-alanine and its effect on muscle carnosine synthesis in human vastus lateralis. *Amino Acids (Vienna)* 30: 279–289, 2006.
19. **Hill CA, Harris RC, Kim HJ, Harris BD, Sale C, Boobis LH, Kim CK, Wise JA.** Influence of beta-alanine supplementation on skeletal muscle carnosine concentrations and high intensity cycling capacity. *Amino Acids (Vienna)* 32: 225–233, 2007.
20. **Horinishi H, Grillo M, Margolis FL.** Purification and characterization of carnosine synthetase from mouse olfactory bulbs. *J Neurochem* 31: 909–919, 1978.
21. **Ibanez J, Pullinen T, Gorostiaga E, Postigo A, Mero A.** Blood lactate and ammonia in short-term anaerobic work following induced alkalosis. *J Sports Med Phys Fitness* 35: 187–193, 1995.
22. **Johnson MA, Polgar J, Weightman D, Appleton D.** Data on the distribution of fibre types in thirty-six human muscles. An autopsy study. *J Neurol Sci* 18: 111–129, 1973.
23. **Kindermann W, Keul J, Huber G.** Physical exercise after induced alkalosis (bicarbonate or tris-buffer). *Eur J Appl Physiol* 37: 197–204, 1977.
24. **Kreis R, Boesch C.** Orientation dependence is the rule, not the exception in 1H-MR spectra of skeletal muscle: the case of carnosine (Online abstract). *ISMRM Proc Intl Soc Magn Reson Med* 2000. <http://cds.ismrm.org/ismrm-2000/START.PDF> [2007].
25. **Lamb GD, Stephenson DG, Bangsbo J, Juel C.** Point:counterpoint: Lactic acid accumulation is an advantage/disadvantage during muscle activity. *J Appl Physiol* 100: 1410–1414, 2006.
26. **Lamont C, Miller DJ.** Calcium sensitizing action of carnosine and other endogenous imidazoles in chemically skinned striated muscle. *J Physiol* 454: 421–434, 1992.
27. **Linderman JK, Gosselink KL.** The effects of sodium bicarbonate ingestion on exercise performance. *Sports Med* 18: 75–80, 1994.
28. **Maynard LM, Boissonneault GA, Chow CK, Bruckner GG.** High levels of dietary carnosine are associated with increased concentrations of carnosine and histidine in rat soleus muscle. *J Nutr* 131: 287–290, 2001.
29. **Messonnier L, Kristensen M, Juel C, Denis C.** Importance of pH regulation and lactate/H<sup>+</sup> transport capacity for work production during supramaximal exercise in humans. *J Appl Physiol* 102: 1936–1944, 2007.
30. **Pan JW, Hamm JR, Rothman DL, Shulman RG.** Intracellular pH in human skeletal muscle by <sup>1</sup>H NMR. *Proc Natl Acad Sci USA* 85: 7836–7839, 1988.
31. **Parkhouse WS, McKenzie DC, Hochachka PW, Ovalle WK.** Buffering capacity of deproteinized human vastus lateralis muscle. *J Appl Physiol* 58: 14–17, 1985.
32. **Ponte J, Harris RC, Hill CA, Sale C, Jones GA, Kim HJ, Wise JA.** Effect of 14 and 28 days  $\beta$ -alanine supplementation on isometric endurance of the knee extensors (Abstract). *J Sports Sci* 25: 344, 2006.
33. **Raymer GH, Marsh GD, Kowalchuk JM, Thompson RT.** Metabolic effects of induced alkalosis during progressive forearm exercise to fatigue. *J Appl Physiol* 96: 2050–2056, 2004.
34. **Requena B, Zabala M, Padial P, Ferliche B.** Sodium bicarbonate and sodium citrate: ergogenic aids? *J Strength Cond Res* 19: 213–224, 2005.
35. **Rohmert W.** Ermittlung von Erholungspausen für statische Arbeit des Menschen. *Int Z Angew Physiol Einschl Arbeitsphysiol* 18: 123–164, 1960.
36. **Schroder L, Bachert P.** Evidence for a dipolar-coupled AM system in carnosine in human calf muscle from in vivo <sup>1</sup>H NMR spectroscopy. *J Magn Reson* 164: 256–269, 2003.
37. **Schroder L, Schmitz C, Bachert P.** Molecular dynamics and information on possible sites of interaction of intramyocellular metabolites in vivo from resolved dipolar couplings in localized <sup>1</sup>H NMR spectra. *J Magn Reson* 171: 213–224, 2004.
38. **Soher BJ, van Zijl PC, Duyn JH, Barker PB.** Quantitative proton MR spectroscopic imaging of the human brain. *Magn Reson Med* 35: 356–363, 1996.
39. **Stout JR, Cramer JT, Zoeller RF, Torok D, Costa P, Hoffman JR, Harris RC.** Effects of beta-alanine supplementation on the onset of neuromuscular fatigue and ventilatory threshold in women. *Amino Acids (Vienna)* 32: 381–386, 2007.
40. **Suyama M, Suzuki T, Maruyama M, Saito K.** Determination of carnosine, anserine and balenine in the muscle of animals. *Bull Jpn Soc Sci Fish* 36: 1048–1053, 1970.
41. **Suzuki Y, Ito O, Mukai N, Takahashi H, Takamatsu K.** High level of skeletal muscle carnosine contributes to the latter half of exercise performance during 30-s maximal cycle ergometer sprinting. *Jpn J Physiol* 52: 199–205, 2002.
42. **Tanokura M, Tasumi M, Miyazawa T.** <sup>1</sup>H nuclear magnetic resonance studies of histidine-containing di- and tripeptides. Estimation of the effects of charged groups on the pKa value of the imidazole ring. *Biopolymers* 15: 393–401, 1976.
43. **Tiryaki GR, Atterbom HA.** The effects of sodium bicarbonate and sodium citrate on 600 m running time of trained females. *J Sports Med Phys Fitness* 35: 194–198, 1995.
44. **Tofts P.** *Quantitative MRI of the Brain: Measuring Changes Caused by Disease.* Chichester, UK: Wiley, 2003.
45. **Wilkes D, Gledhill N, Smyth R.** Effect of acute induced metabolic alkalosis on 800-m racing time. *Med Sci Sports Exerc* 15: 277–280, 1983.
46. **Yoshioka Y, Oikawa H, Ehara S, Inoue T, Ogawa A, Kanbara Y, Kubokawa M.** Noninvasive measurement of temperature and fractional dissociation of imidazole in human lower leg muscles using 1H-nuclear magnetic resonance spectroscopy. *J Appl Physiol* 98: 282–287, 2005.