Articles in PresS. J Appl Physiol (August 5, 2010). doi:10.1152/japplphysiol.00563.2010

The Two-Hour Marathon: Who and When?

M.J. Joyner*

J.R. Ruiz†

A. Lucia§

*Department of Anesthesiology
Mayo Clinic
Rochester, MN

†Department of Biosciences and Nutrition
Unit for Preventive Nutrition
Karolinska Institutet
Stockholm, Sweden

§Universidad Europea de Madrid
Spain

Corresponding Author: Michael J. Joyner, M.D.
Department of Anesthesiology
Mayo Clinic
200 First Street SW
Rochester, MN 55905

Phone (507) 255-4288
Fax (507) 255-7300
E-mail: joyner.michael@mayo.edu

Copyright © 2010 by the American Physiological Society.

OVERVIEW

In this Viewpoint we ask if information about the physiology, genetics, and empirical history of elite endurance performance can provide insight into the question of "who" will break the two-hour marathon barrier and when this might happen. We also identify several physiological questions that we believe need attention.

The current world record in the men's marathon is 2:03:58 (Gebrselassie 2007). This record has fallen by more than 16 minutes since the early 1950s after high volume/year round training was adopted widely. Except for the 1970s, the record has fallen by ~1-5 minutes per decade since 1960 when Africans entered international competition. Improvements since 1980 likely also reflect increased prize money and competitive opportunities that allowed top athletes to earn a living running. Figure 1 shows the history of marathon times and projected improvements. Using times from 1960, the open squares suggest it will take 12-13 years to break 2 hours assuming a ~20 sec reduction per year. If times from 1980 are used the filled squares suggest it will take 25 years assuming a ~10 sec reduction per year. Consistent with the idea that marked improvement is likely, empirical models of running times suggest that the men's world records for the 10,000m and half marathon are equivalent to a marathon time of ~2:02 - 2:03 (5,21).

Physiology of the Two-Hour Marathon

The physiological determinants of distance running performance (VO_2max, lactate threshold, and running economy) have been used to develop a model of marathon

65 performance (9,10). Elite marathon runners typically have VO_2 max values ranging
66 from ~70 ml/kg/min to ~85 ml/kg/min. These individuals can sustain running speeds
67 that require 85-90% VO_2 max for more than one hour, and these factors along with
68 knowledge of the oxygen cost to run a given speed (running economy) provide a
69 reasonable estimate of marathon pace (9,10). When outstanding values for these three
70 key variables are used in this model, a sub- two hour marathon seems physiologically
71 possible.
72
73 While there are many possible combinations that might lead to elite performances, it
74 appears that extremely high values for VO_2max and outstanding running economy are
75 rarely seen in the same person (9,10). East African runners do not have particularly
76 exceptional values for VO_2max or lactate threshold, but generally have outstanding
77 running economy (13,14,23). The classic study of Pollock showed that elite distance
78 runners who focused on the marathon had lower VO_2max values and better running
79 economy that those who focused on shorter races (19). Based on these data and other
80 anecdotal reports, it appears that whoever breaks two hours for the marathon will have
81 exceptional running economy (2, 4).
82
83 In this context, there is clearly a need for more information about the relationship
84 between VO_2max and running economy and the physiological explanation for the
85 relationship *if it exists.* There is evidence that VO_2max and gross mechanical efficiency
86 are inversely related in cyclists and influenced by muscle fiber type (16). By contrast,
87 running economy seems more related to mechanical factors including vertical

displacement and so-called braking on foot strike (11,24). Exceptional running economy might also provide two important physiological advantages. First, fuel utilization would be lower and perhaps glycogen depletion delayed. Second, metabolic heat production would also be lower potentially reducing thermal stress. To our knowledge these potential advantages have not be studied extensively.

What will the Two-Hour Marathoner Look Like?

Forty-one of the 50 fastest marathons have been run by Kenyans or Ethiopians (1). Importantly, the mean height and weight of the 30 runners (29 Africans) who have broken 27 minutes for 10,000 m is `170± 6 cm, and 56±5 kg, with only one runner greater than 178 cm or 70 kg (12). Additionally, most of these athletes had exposure to high altitude and significant physical activity early in life. In this context, small body size has a favorable effect on VO_2 max; however, less is known about its influence on running economy (7).

From these observations other questions emerge: (i) Does exposure to the combination of high altitude and physical activity early in life lead to pulmonary adaptations that reduce the incidence of arterial desaturation seen during heavy exercise in elite athletes (3,5,15,16)? and (ii) would the reduction in metabolic heat production along with a favorable body weight to surface area ratio have the net affect of reducing thermoregulatory stress during periods of prolonged, intense exercise? While these questions might be difficult to study, small differences could be decisive when races are won and records set by very small margins. However, there are examples of “big”

111 runners like Paula Radcliffe, Ron Clarke and Derek Clayton who have been highly
112 successful. Importantly, Radcliffe and Clayton are known to have superb running
113 economy, and Radcliffe's running economy improved dramatically over time, providing
114 at least some evidence that this factor is "traininable" (8,19).
115
116 Genotype: Probabilistic versus Deterministic
117 Genetic factors may limit or enhance the possibility of running a very fast marathon. At
118 present much of what is known comes from association studies, with the angiotensin
119 converting enzyme (*ACE*) I/D and α-actinin-3 (*ACTN3*) R577X gene polymorphisms
120 having been studied extensively. The *ACE* I allele is theoretically associated with
121 improved cardiovascular function during exercise, and could also favor muscle
122 efficiency (26). While there is an overrepresentation of the I allele in the best Spanish
123 marathon runners (sub 2:09 marathon performance) (15), the *ACE* I/D polymorphism is
124 not associated with the success of the best elite endurance runners worldwide,
125 including Kenyans (25). The association between the *ACTN3* R577X variation and elite
126 'power 'athlete status is strongly documented (27), yet this is not the case for endurance
127 running (28).
128
129 Beyond potential genotype/phenotype associations (which are yet to be clearly
130 established in elite marathoners), the task of quantifying the genetic contribution to elite
131 marathon performance is challenging. A record holders's phenotype results from the
132 combined influence of hundreds of genes, epigenetic factors, and non-hereditary
133 environmental influences. Using algorithms that take into account the combined

134 influence of several candidate gene variants associated with endurance performance
135 [i.e., the so-called 'total genotype score' (TGS), ranging from 0 to 100], it appears that
136 genetic factors increases the possibility of becoming a marathon champion (22). For
137 example, a Caucasian individual with a TGS value above 75 has ~5 times greater
138 chance of achieving elite endurance runner status compared to those with a TGS below
139 75. Yet, less than half of the best Spanish marathoners have TGS values above 75;
140 and, using this approach it is estimated there are nearly 6 million Spanish individuals
141 with the 'genetic' potential for elite marathon performance. Whether having the best
142 possible TGS (i.e. 100) increases the odds of breaking two-hours is unknown.
143
144 Summary
145 Whoever breaks two hours will likely have outstanding running economy and small body
146 size along with exposure to high altitude, and significant physical activity early in life.
147 However, neither these factors nor any specific suite of genotypes appear to be
148 obligatory for a time this fast. Current trends suggest that an East African will be the
149 first to break two hours. However periods of regional dominance in distance running are
150 not unique to the East Africans: athletes from Finland, Eastern Europe, Australia and
151 New Zealand have all had extended periods of success at a range of distances (17).
152 From a physiological perspective, more information is clearly needed on the relationship
153 between VO_2max and running economy and the influence of running economy and
154 body size on thermoregulation and fuel use.
155

References

1. **Association of Road Racing Statisticians.** www.arrs.net.

2. **Billat V, Lepretre PM, Heugas AM, Laurence MH, Salim D, Koralsztein JPA.** Training and bioenergetic characteristics in elite male and female Kenyan runners. *Med Sci Sports Exerc* 35: 297-304, 2003.

3. **Dempsey JA, Hanson PG, Henderson KS.** Exercise-induced arterial hypoxaemia in healthy human subjects at sea level. *J Physiol* 355: 161-175, 1984.

4. **Foster C, Lucia A.** Running economy: the forgotten factor in elite performance. *Sports Med* 37: 316-319, 2007.

5. **Gardner JB, Purdy JG**, Computerized Running Training Programs. TAFNEWS PRESS, Los Altos, CO 1970

6. **Harms CA, McClaran SR, Nickele GA, Pegelow DR, Nelson WB, Dempsey JA.** Exercise-induced arterial hypoxaemia in healthy young women. *J Physiol* 507: 619-628, 1998.

7. **Jensen K, Johansen L, Secher NH.** Influence of body mass on maximal oxygen uptake: effect of sample size. *Eur J Appl Physiol* 84: 201-5, 2001.

8. **Jones AM.** A five year physiological case study of an Olympic runner. *Br J Sports Med* 32: 39-43, 1998.

9. **Joyner MJ**. Modeling: optimal marathon performance on the basis of physiological factors. *J Appl Physiol* 70: 683-687, 1991.

10. **Joyner MJ, Coyle EF.** Endurance exercise performance: the physiology of champions. *J Physiol* 586: 35-44, 2007.

11. **Kyröläinen H, Belli A, Komi PV.** Biomechanical factors affecting running economy. *Med Sci Sports Exerc* 33: 1330-1337, 2001.

12. **LetsRun.com.** Chris Solinsky is the Tallest and Heaviest Sub-27 man in History. www.letsrun.com/2010/heightweight0504.php.

13. **Larsen HB.** Kenyan dominance in distance running. *Comp Biochem Physiol A Molec Integr Physiol* 136: 161-170, 2003.

14. **Lucia A, Esteve-Lanao J, Oliván J, Gómez-Gallego F, San Juan AF, Santiago C, Pérez M, Chamorro-Viña C, Foster C.** Physiological

201 characteristics of the best Eritrean runners-exceptional running economy. *Appl*
202 *Physiol Nutr Metab* 31: 530-540, 2006.
203
204 15. **Lucía A, Gómez-Gallego F, Chicharro JL, Hoyos J, Celaya K, Córdova A,**
205 **Villa G, Alonso JM, Barriopedro M, Pérez M, Earnest CP.** Is there an
206 association between ACE and CKMM polymorphisms and cycling performance
207 status during 3-week races*?* *Int J Sports Med* 26: 442-447, 2005.
208
209 16. **Lucia A, Hoyos J, Pérez M, Santalla A, Chicharro JL.** Inverse relationship
210 between VO_2max and economy/efficiency in world-class cyclists. Med Sci Sports
211 Exerc 34: 2079–2084, 2002.
212
213 17. **Megede EZ, Hymans R.** *Progression of World Best Performances and IAAF*
214 *Approved World Records.* Monaco: International Athletic Foundation, 1991, 705
215 pp.
216
217 18. **Nielsen HB.** Arterial desaturation during exercise in man: implication for O2
218 uptake and work capacity. *Scand J Med Sci Sports* 13: 339-358, 2003.
219
220 19. **Pollock ML.** Submaximal and maximal working capacity of elite distance
221 runners. Part I: Cardiorespiratory aspects. *Ann NY Acad Sci* 301: 310-322,
222 1977.
223
224 20. **Ravikumar P, Bellotto DJ, Johnson RL Jr, Hsia CC.** Permanent alveolar
225 remodeling in canine lung induced by high-altitude residence during maturation.
226 *J Appl Physiol* 107: 1911-1917, 2009.
227
228 21. **Riegel PS.** Athletic records and human endurance. *Am Scientist* 69: 285-290,
229 1981.
230
231 22. **Ruiz JR, Gómez-Gallego F, Santiago C, González-Freire M, Verde Z, Foster**
232 **C, Lucia A.** Is there an optimum endurance polygenic profile? *J Physiol* 587:
233 1527-1534, 2009.
234
235 23. **Saltin B, Larsen H, Terrados N, Bangsbo J, Bak T, Kim CK, Svedenhag J,**
236 **Rolf CJ.** Aerobic exercise capacity at sea level and at altitude in Kenyan boys,
237 junior and senior runners compared with Scandinavian runners. *Scand J Med*
238 *Sci Sports* 5: 209-221, 1995.
239
240 24. **Saunders PU, Pyne DB, Telford RD, Hawley JA.** Factors affecting running
241 economy in trained distance runners. *Sports Med* 34: 465-485, 2004.
242
243 25. **Scott RA, Moran C, Wilson RH, Onywera V, Boit MK, Goodwin WH, Gohlke**
244 **P, Payne J, Montgomery H, Pitsiladis YP.** No association between angiotensin
245 converting enzyme (ACE) gene variation and endurance athlete status in
246 Kenyans. *Comp Biochem Physiol A Mol Integr Physiol* 141: 169-175, 2005.

26. **Williams AG, Rayson MP, Jubb M, World M, Woods DR, Hayward M, Martin J, Humphries SE, Montgomery HE.** The ACE gene and muscle performance. *Nature* 403:614, 2000.

27. **Yang N, MacArthur DG, Gulbin JP, Hahn AG, Beggs AH, Easteal SH, North K.** ACTN3 genotype is associated with human elite athletic performance. *Am J Hum Genet* 73: 627-631, 2003.

28. **Yang N, MacArthur DG, Wolde B, Onywera VO, Boit MK, Lau SY, Wilson RH, Scott RA, Pitsiladis YP, North K.** The ACTN3 R577X polymorphism in East and West African athletes. *Med Sci Sports Exerc* 39: 1985-1988 2007.

Figure Legend

Figure 1. Progression of world record times in the marathon since the late 1920s. The rapid fall in record time in the 50s and 60s likely reflects: i) the widespread adoption of high volume/year round training after WWII; and ii) the participation of East-African runners in international competition starting in the 1960s. There was limited progress during the 1970s, but the record has fallen more than 5 minutes over the last ~30 years. On average, there has been ~20 s reduction per year since 1960. The open squares show that if this rate of improvement continues, a time under 2 hours could occur in 12-13 years (by 2021-2022). The closed squares show that if only data from 1980 are used, a time under 2 hours would occur in ~25 years based on an estimated improvement of ~10s per year. The recent increase in the number of high profile races on fast courses that offer substantial prize money may also contribute to faster world records in the near future.


Time (minutes)
152
150
148
146
144
142
140
138
136
134
132
130
128
126
124
122
120
118
1925 1930 1935 1940 1945 1950 1955 1960 1965 1970 1975 1980 1985 1990 1995 2000 2005 2010 2015 2020 2025 2030 2035 2040
Year